

# Biological Rhythm and Dynamics through Chemistry

## Research Center of Integrative Molecular Systems Division of Trans-Hierarchical Molecular Systems



**AKIYAMA, Shuji**  
Professor  
[akiyamas@ims.ac.jp]

### Education

1997 B.E. Kyoto University  
1999 M.E. Kyoto University  
2002 Ph.D. Kyoto University

### Professional Employment

2001 JSPS Research Fellow  
2002 JSPS Postdoctoral Fellow  
2003 RIKEN Special Postdoctoral Researcher  
2005 JST-PRESTO Researcher  
2008 Junior Associate Professor, Nagoya University  
2011 Associate Professor, Nagoya University  
2012 Professor, Institute for Molecular Science  
Professor, The Graduate University for Advanced Studies

### Awards

2016 The 13<sup>th</sup> (FY2016) JSPS PRIZE  
2008 The Commendation for Science and Technology by the Minister of Education, Culture, Sports, Science and Technology  
The Young Scientists' Prize  
2007 Young Scientist Prize, The Biophysical Society of Japan  
2006 SAS Young Scientist Prize, IUCr Commission on Small-angle Scattering  
2002 The Protein Society Annual Poster Board Award

### Member

Assistant Professor  
MUKAIYAMA, Atsushi  
FURUIKE, Yoshihiko  
Post-Doctoral Fellow  
OUYANG, Dongyan  
Visiting Scientist  
TONIOLO, Paul\*  
MARIE, Vincent\*  
Graduate Student  
SIMON, Damien  
Technical Fellow  
SHINTANI, Atsuko  
HAYAKAWA, Arina  
WASHIO, Midori  
TAKAYA, Mari  
SUGISAKA, Kanae  
Secretary  
SUZUKI, Hiroko

**Keywords** Biological Rhythm, Circadian Clock, Cyanobacteria

Living organisms on Earth evolved over time to adapt to daily environmental alterations, and eventually acquired endogenous time-measuring (biological clock) systems. Various daily activities that we perform subconsciously are controlled by the biological clock systems sharing three characteristics. First, the autonomic rhythm repeats with an approximately 24-hour (circadian) cycle (self-sustainment). Second, the period is unaffected by temperature (temperature compensation). Third, the phase of the clock is synchronized with that of the outer world in response to external stimuli (synchronization). We seek to explain these three characteristics, and consider the biological clock system of cyanobacteria to be an ideal experimental model.

The major reason that cyanobacteria are considered to be the ideal experimental model is that the core oscillator that possesses the three characteristics of the clock can be easily reconstructed within a test tube. When mixing the three clock proteins KaiA, KaiB, and KaiC with ATP, the structure and enzyme activity of KaiC change rhythmically during a circadian cycle. Taking advantage of this test tube experiment, we used an approach combining biology, chemistry, and physics

to elucidate the means by which the clock system extends from the cellular to atomic levels.

Among the three Kai proteins, KaiC is the core protein of the oscillator. In the presence of KaiA and KaiB, KaiC reveals the rhythm of autophosphorylation and dephosphorylation; however, the cycle of this rhythm depends on the ATPase activity of KaiC independent of KaiA or KaiB. For example, when the ATPase activity of KaiC doubles as a result of amino acid mutations, the frequencies of both the *in vitro* oscillator and the intracellular rhythm also double (the cycle period is reduced to half). This mysterious characteristic is called a transmembrane hierarchy, in which the cycle (frequency) and even the temperature compensation both *in vitro* and *in vivo* are greatly affected (controlled) by the function and structure of KaiC.


How are the circadian activities and temperature compensation features encoded in KaiC and then decoded from it to propagate rhythms at the cellular level? We are committed to better understanding biological clocks and other dynamic systems through the chemistry of circadian **rhythm, structure**, and evolutionary **diversity**.

### Selected Publications

- D. Ouyang, Y. Furuike, A. Mukaiyama, K. Ito-Miwa, T. Kondo and S. Akiyama, *Int. J. Mol. Sci.* **20**, 2789–2800 (2019).
- A. Mukaiyama, D. Ouyang, Y. Furuike and S. Akiyama, *Int. J. Biol. Macromol.* **131**, 67–73 (2019).
- A. Mukaiyama, Y. Furuike, J. Abe, E. Yamashita, T. Kondo and S. Akiyama, *Sci. Rep.* **8**, 8803 (2018).
- J. Abe, T. B. Hiyama, A. Mukaiyama, S. Son, T. Mori, S. Saito, M. Osako, J. Wolanin, E. Yamashita, T. Kondo and S. Akiyama, *Science* **349**, 312–316 (2015).
- Y. Murayama, A. Mukaiyama, K. Imai, Y. Onoue, A. Tsunoda, A. Nohara, T. Ishida, Y. Maéda, T. Kondo and S. Akiyama, *EMBO J.* **30**, 68–78 (2011).
- S. Akiyama, A. Nohara, K. Ito and Y. Maéda, *Mol. Cell* **29**, 703–716 (2008).

## 1. Structure: Atomic-Scale Origins of Clock Slowness in Cyanobacterial Circadian Clock System<sup>1)</sup>


To identify the structural origins of slowness encoded in KaiC, its N-terminal ATPase domain was analyzed using high-resolution x-ray crystallography.<sup>1)</sup> Water molecules are prevented from attacking into the ideal position (a black dot in Figure 1) for the ATP hydrolysis by a steric hindrance near ATP phosphoryl groups. In addition, this hindrance is surely anchored to a spring-like structure derived from polypeptide isomerization. The ATP hydrolysis, which involves access of a water molecule to the bound ATP and reverse isomerization of the polypeptide, is expected to require a significantly larger amount of free energy than for typical ATP hydrolysis. The atomic structure discovered by us explains why the ATPase activity of KaiC is so much lower (by 100- to 1,000,000-fold) than that of typical ATPase molecules.



**Figure 1.** Structural basis for steady slowness. The steric barrier prevents access of a water molecule to the catalytic site (indicated by a black dot).

## 2. Rhythm: Transmural Hierarchy in Cyanobacterial Circadian Clock System<sup>2,3)</sup>

KaiC ATPase is of particular interest here, as it finely correlates to the frequencies of *in vivo* as well as *in vitro* oscillations. This unique property has inspired us to develop an ATPase-based screening for KaiC clock mutants giving short, long, and/or temperature-dependent periods.<sup>2)</sup> A developed HPLC system with a 4-channel temperature controller has reduced approximately 80% of time costs for the overall screening process (Figure 2).




**Figure 2.** Development of a quick ATPase assay system.

How is the intra-molecular slowness encoded in KaiC (Figure 1) transmitted to the inter-molecular interactions with other Kai proteins? To address this question, a tryptophan residue was introduced in the N-terminal ring of KaiC as the fluorescent probe for KaiBC complex formation.<sup>3)</sup> Our kinetic data indicated that KaiB exclusively selects the post-ATP-hydrolysis state of KaiC to form the KaiBC complex. This process follows a mechanism called conformational selection (CS), in which proteins (KaiC) first undergoes a structural change to form a specific intermediate. Ligands (KaiB) are

then recognized specifically through the intermediate state to form a tight ligand-protein complex. The CS mechanism is elegantly designed in KaiC so that the slow intra-molecular ATPase reaction in KaiC can be the rate-limiting step of the overall KaiBC complex formation.

## 3. beyond Evolutionary Diversity<sup>1,4)</sup>

In the presence of KaiA and KaiB, the ATPase activity of KaiC oscillates on a 24-hour cycle. KaiC is not capable of maintaining a stable rhythm on its own, but its activity was observed to fluctuate with reduced amplitude over time (Figure 3A). We have identified a signal component that is similar to damped oscillation, and propose that it encodes the specific frequency, equivalent to a 24-hour cycle.<sup>1)</sup>



**Figure 3.** Damped oscillation of KaiC ATPase activity (A) and evolutionary diversity of cyanobacteria (B).

As this proposal is further discussed and verified, development of new studies is expected. Some KaiA and KaiB genes are missing in several strains of cyanobacteria. This is understandable to some extent if KaiC possesses the specific frequency. Given our current understanding of this phenomenon, *what specific frequencies are possessed by KaiCs and their homologues in other species and ancestral cyanobacteria?* (Figure 3B) The habitats of cyanobacteria are diverse, so the space of their sequence is immense.<sup>4)</sup> If you strain your ears, the rhythms of the ancient Earth may be heard from beyond evolutionary diversity.

## 4. Bio-SAXS Activity in IMS<sup>5,6)</sup>

We have supported SAXS users so that they can complete experiments smoothly and publish their results.<sup>5,6)</sup>

### References

- 1) J. Abe, T. B. Hiyama, A. Mukaiyama, S. Son, T. Mori, S. Saito, M. Osako, J. Wolanin, E. Yamashita, T. Kondo and S. Akiyama, *Science* **349**, 312–316 (2015).
- 2) D. Ouyang, Y. Furuike, A. Mukaiyama, K. Ito-Miwa, T. Kondo and S. Akiyama, *Int. J. Mol. Sci.* **20**, 2789–2800 (2019).
- 3) A. Mukaiyama, Y. Furuike, J. Abe, E. Yamashita, T. Kondo and S. Akiyama, *Sci. Rep.* **8**, 8803 (2018).
- 4) A. Mukaiyama, D. Ouyang, Y. Furuike and S. Akiyama, *Int. J. Biol. Macromol.* **131**, 67–73 (2019).
- 5) M. Okumura, K. Noi, S. Kanemura, M. Kinoshita, T. Saio, Y. Inoue, T. Hikima, S. Akiyama, Y. Ogura and K. Inaba, *Nat. Chem. Biol.* **15**, 499–509 (2019).
- 6) Y. Furukawa, C. Lim, T. Tosha, K. Yoshida, T. Hagai, S. Akiyama, S. Watanabe, K. Nakagome and Y. Shiro, *PLoS One* **13**, e0204355 (2018).

\* IMS International Internship Program