

Angle-Resolved Photoemission Study on Strongly Correlated Electron Materials

UVSOR Synchrotron Facility Division of Advanced Solid State Physics



TANAKA, Kiyohisa
Associate Professor
[k-tanaka@ims.ac.jp]

Education

2000 B.S. The University of Tokyo
2005 Ph.D. The University of Tokyo

Professional Employment

2005 Postdoctoral Fellow, Stanford University and Lawrence Berkeley National Laboratory
2008 Assistant Professor, Osaka University
2013 Associate Professor, Osaka University
2014 Associate Professor, Institute for Molecular Science
Associate Professor, The Graduate University for Advanced Studies

Member

Graduate Student
HOSHINA, Takumi*
SUGIMOTO, Takafumi*
KOYAMA, Shotaro*
MITA, Manaya*

Secretary

ISHIHARA, Mayumi
KAMO, Kyoko
YOKOTA, Mitsuyo


Keywords

Strongly Correlated Electron System, Synchrotron Light, Photoemission

Strongly correlated electron materials have attracted more attentions in the last few decades because of their unusual and fascinating properties such as high- T_c superconductivity, giant magnetoresistance, heavy fermion and so on. Those unique properties can offer a route toward the next-generation devices. We investigate the mechanism of the physical properties as well as the electronic structure of those materials by using angle-resolved photoemission spectroscopy (ARPES). ARPES is a powerful experimental technique, directly measuring the energy (E) and momentum (k) relation, namely the band structure of solids. In the last quarter of a century, the energy resolution and angular resolution of ARPES have improved almost three order of magnitude better, which makes us possible to address the fine structure of the electronic structure near the Fermi level: Superconducting gap, kink structure and so on. The main target materials of our group is high- T_c superconductors, such as cuprates and iron pnictides and use UVSOR-III as a strong light source.

Our group is also developing high-efficiency spin-resolved ARPES system. Spintronics is a rapidly emerging field of science and technology that will most likely have a significant

impact on the future of all aspects of electronics as we continue to move into the 21st century. Understanding magnetism of surfaces, interfaces, and nanostructures is greatly important for realizing the spintronics which aims to control and use the function of spin as well as the charge of electrons. Spin-resolved ARPES is one of the most powerful experimental techniques to investigate the magnetic properties of such materials.



Selected Publications

- K. Tanaka, W. S. Lee, D. H. Lu, A. Fujimori, T. Fujii, Risdiana, I. Terasaki, D. J. Scalapino, T. P. Devereaux, Z. Hussain and Z.-X. Shen, "Distinct Fermi-Momentum-Dependent Energy Gaps in Deeply Underdoped Bi2212," *Science* **314**, 1910–1913 (2006).
- W. S. Lee, I. M. Vishik, K. Tanaka, D. H. Lu, T. Sasagawa, N. Nagaosa, T. P. Devereaux, Z. Hussain and Z.-X. Shen, "Abrupt Onset of a Second Energy Gap at the Superconducting Transition of Underdoped Bi2212," *Nature* **450**, 81–84 (2007).
- K. Tanaka, N. Hieu, G. Vincini, T. Masui, S. Miyasaka, S. Tajima and T. Sasagawa, "Quantitative Comparison between Electronic Raman Scattering and Angle-Resolved Photoemission Spectra in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ Superconductors: Doping Dependence of Nodal and Antinodal Superconducting Gaps," *J. Phys. Soc. Jpn.* **88**, 044710 (2019).
- S. Ideta, N. Murai, M. Nakajima, R. Kajimoto and K. Tanaka, "Experimental Investigation of the Suppressed Superconducting Gap and Double-Resonance Mode in $\text{Ba}_{1-x}\text{K}_x\text{Fe}_2\text{As}_2$," *Phys. Rev. B* **100**, 235135 (7 pages) (2019).

1. Development of Spin-Resolved ARPES with Image-Spin Detection

Spintronics is a rapidly emerging field of science and technology that will most likely have a significant impact on the future of all aspects of electronics as we continue to move into the 21st century. Understanding magnetism of surfaces, interfaces, and nanostructures is greatly important for realizing the spintronics which aims to control and use the function of spin as well as the charge of electrons. Spin- and angle-resolved photoemission spectroscopy (spin-resolved ARPES) is one of the most powerful experimental techniques to investigate the magnetic properties of such materials, where one can know the “complete” information of the electronic states of materials; energy, momentum, and spin direction. Recent development of high energy and angle resolved photoelectron analyzer as well as the contemporary light sources such as third generation synchrotron radiation make it possible for the photoemission spectroscopy to investigate not only band structures but many body interactions of electrons in solids. However, appending the spin resolution to photoemission spectroscopy is quite difficult because of an extremely low efficiency (10^{-4}) of Mott-type spin detectors. Recently, very-low-energy-electron-diffraction (VLEED-type) spin detector with 100 times higher efficiency than that of conventional Mott-type has been developed and spin-resolved ARPES has been started to be realized. So far, most of the spin-resolved ARPES systems in the world are using the single-channel detector and efficiency is still a problem.

Beamline BL5U at UVSOR has been totally reconstructed by our group, and opened for users as high photon flux and high energy resolution ARPES beamline since 2017. As a new function for this beamline, we have started high-efficient spin-resolved ARPES project with multi-channel detection (we call “image-spin” detection). The goal of this project is to realize the 100 times better efficiency and the 10 times better momentum resolution than the current spin-resolved ARPES system in the world, which can be a breakthrough in this field.

In 2020, we set up the spin detection apparatus shown in Figure 1(a) and finished tuning the electron lens parameters of the spin detection section. With this apparatus, we succeeded in obtaining a spin-resolved signal on the Au(111) surface as shown in Figure 2(c). According to rough estimates, the




Figure 1. Previous (a) and current (b) setup of image-spin ARPES.

efficiency of spin-resolved ARPES was 100 times higher than that of the single-channel detection systems currently used in the world. However, the spin-resolved ARPES bands were broad compared to the normal ARPES ones shown in Figure 2(b), meaning that the momentum resolution was not so good. With this apparatus, the instrument required two VLEED detection chambers and a high-quality spin target of the same quality to detect spin information in the x , y , and z directions of the sample. In addition, the spin targets had to be magnetized frequently during the measurement.

To overcome these problems, we have introduced a new “spin manipulator” that can change the spin direction of the passing electrons in any direction. The new setup shown in Figure 1(b) requires only one VLEED detection chamber and the users do not need to magnetize the spin target during the measurement. The installation of the spin manipulator and optimization of the spin target deposition conditions have greatly improved the spin-resolved images, and in 2022, we were able to obtain spin-resolved images with momentum resolution comparable to that of normal ARPES, as shown in Figure 2(d). We are currently optimizing the lens parameters of the spin manipulator to obtain spin information in the remaining two axial directions.




Figure 2. a) Fermi surface of Rashba spin splitting in Au(111) surface states and (b) image plot of normal ARPES along the yellow bar in (a). Spin-resolved ARPES data showing the spin polarization (blue–red scale) with the previous experimental setup (c) and the current setup (d).