

# Development of High-Precision Coherent Control and Its Applications

Department of Photo-Molecular Science  
Division of Photo-Molecular Science II




OHMORI, Kenji	Professor
KATSUKI, Hiroyuki	Assistant Professor
HOSAKA, Kouichi	IMS Fellow*
DELAGNES, Jean-Christophe	Post-Doctoral Fellow†
SHIMADA, Hiroyuki	Post-Doctoral Fellow
GOTO, Haruka	Graduate Student
INAGAKI, Itsuko	Secretary

Coherent control is based on manipulation of quantum phases of wave functions. It is a basic scheme of controlling a variety of quantum systems from simple atoms to nanostructures with possible applications to novel quantum technologies such as bond-selective chemistry and quantum computation. Coherent control is thus currently one of the principal subjects of various fields of science and technology such as atomic and molecular physics, solid-state physics, quantum electronics, and information science and technology. One promising strategy to carry out coherent control is to use coherent light to modulate a matter wave with its optical phase. We have so far developed a high-precision wave-packet interferometry by stabilizing the relative quantum phase of the two molecular wave packets generated by a pair of femtosecond laser pulses on the attosecond time scale. We will apply our high-precision quantum interferometry to gas, liquid, solid, and surface systems to explore and control various quantum phenomena.


## 1. Actively Tailored Spatiotemporal Images of Quantum Interference on the Picometer and Femtosecond Scales<sup>1)</sup>

Interference fringes of quantum waves weave highly regular space-time images, which could be seen in various wave systems such as wave packets in atoms and molecules, Bose Einstein condensates, and Fermions in a box potential. We have experimentally designed and visualized spatiotemporal images of dynamical quantum interferences of two counter-

propagating nuclear wave packets in the iodine molecule; the wave packets are generated with a pair of femtosecond laser pulses whose relative phase is locked within the attosecond time scale. The design of the image has picometer and femtosecond resolutions, and changes drastically as we change the relative phase of the laser pulses, providing a direct spatiotemporal control of quantum interferences.



**Figure 1.** Pump-control-probe scheme for tailoring and visualizing the quantum carpet in the iodine molecule. (Left) Three femtosecond laser pulses are employed. Two of them are phase-locked and used as the pump and control pulses, and the other one is not phase-locked and is used as the probe pulse. The fluorescence signal induced by the probe pulse is measured with a photomultiplier tube attached to a monochromator. (Right) The model simulation of the quantum interference of two wave packets created on the *B*-state potential curve of the iodine molecule by the pump and control pulses. The interference shows a highly regular space-time image that looks like a carpet.



**Figure 2.** Comparisons of the quantum carpets measured (left) and simulated (right) at the pump-control relative phases  $\theta_{pc} =$  (a)  $0^\circ$ , (b)  $90^\circ$ , (c)  $180^\circ$ , and (d)  $270^\circ$ . The color scaling is common within each set of measured or simulated carpets; the maxima of those two sets

have the same color. The origin  $\tau_{probe} = 0$  of the simulated carpet denotes a position of the top of the first oscillation around the outer turning point. The simulations include the interactions with the pump, control, and probe pulses. The parameter  $\tau_{control}$ , used for the simulation of  $\theta_{pc} = 0^\circ$ , was 468.920 fs.

#### Reference

- 1) H. Katsuki, H. Chiba, C. Meier, B. Girard and K. Ohmori, *Phys. Rev. Lett.* submitted.

\* Present Affiliation; Postdoctoral Fellow, Japan Atomic Energy Agency

† Present Affiliation; Senior Lecturer, The University of Bordeaux, France